arXiv:1411.0002v1 [math.HO] 31 Oct 2014

A Quick Empirical Reproof of the Asymptotic Normality of the Hirsch Citation Index (First proved by Canfield, Corteel and Savage)

Shalosh B. EKHAD and Doron ZEILBERGER

Abstract: Inspired by Alexander Yong's recent critique of the Hirsch Citation index, we give an empirical (yet **very convincing!**) reproof of the *asymptotic normality* of the Hirsch citation index (alias size of Durfee square) with respect to the uniform distribution on the "sample space" of integer partitions of n. This result was first proved rigorously (but with much greater effort!) by the humans Rodney Canfield, Sylvie Corteel, and Carla Savage. In particular, we confirm the Canfield-Corteel-Savage rigorous evaluation of the average: $(\frac{\sqrt{6}\log 2}{\pi})\sqrt{n}+O(1) = 0.5404446\sqrt{n}+O(1)$, and estimate the variance, numerically, as $0.0811\sqrt{n}+O(1)$, and get estimates extremely close to those of the standard Normal Distribution for the first 12 standardized moments. We also observe that what Yong calls "Rodney Canfield's concentration conjecture", that asserts that most of the "mass" is close to the average, follows immediately from the Canfield-Corteel-Savage 1998 result, since the *variance* is proportional to the average (with a rather small proportionality constant, namely 0.0811/0.5404446, that is approximately 0.1501). All the results in this article were obtained via straightforward symbol- and number- crunching, by the aid of a Maple package called `HIRSCH`, that is available free of charge from `http://www.math.rutgers.edu/~zeilberg/tokhniot/HIRSCH` .

The Cinderella Story of the Size of the Durfee Square

Once upon a time there was an esoteric and specialized notion, called "size of the Durfee square", of interest to at most 100 specialists in the whole world. Then it was kissed by a prince called Jorge Hirsch ([H]), and became the *famous* (and to quite a few people, *infamous*) h-index, of interest to *every* scientist, and scholar, since it tells you how productive a scientist (or scholar) you are!

When Rodney Canfield, Sylvie Corteel, and Carla Savage wrote their beautiful article, [CCS], proving, rigorously, by a very deep and intricate analysis, the *asymptotic normality* of the random variable "size of Durfee square" defined on integer-partitions of n (as $n \to \infty$), with precise asymptotics for the mean and variance, they did not dream that one day their result should be of interest to everyone who has ever published a paper.

Alexander Yong's Critique of the h-index

In the latest issue of the *Notices of the American Mathematical Society*, Alexander Yong contributed ([Y]) a very insightful critique of the Hirsch citation index([H]). Yong mentioned [CCS], but apparently missed its full significance. In particular, what Yong calls "Canfield's concentration conjecture" is an immediate consequence of the fact, proved in [CCS], that what now is called the h-index is asymptotically normal, and the fact, also proved there, that the asymptotic variance is proportional to the asymptotic average.

Why this Redux?

To be honest, we don't have the patience to follow the intricate analysis of [CCS], and while we trust them completely, it is nice to find out things by ourselves. More importantly, we want to describe, via this *case-study*, how one can get much quicker, the same *mathematical knowledge*, by combining *number-crunching* and naive *symbol-crunching* to get (empirically, but very reliably) limiting distributions of many families of combinatorial 'statistics' (or *random-variables*). Often this method can be used to get fully rigorous results, (see [Z1][Z2]), but with much lesser effort, one can get empirical proofs. It is also very easy to come up with examples where a fully rigorous proof is completely beyond the scope of humans, or even computers, and when it is, it is not worth the efforts!

What is the h-index (alias Durfee Square)

Recall that the size of the Durfee square, alias h-index, is defined as follows. For a partition of a positive integer, n, $\lambda = (\lambda_1, \ldots, \lambda_k)$, (where $\lambda_1 \geq \ldots \geq \lambda_k \geq 1$, and $\lambda_1 + \ldots + \lambda_k = n$), $h(\lambda)$ is the largest i such that $\lambda_i \geq i$.

In this note we reprove the above-mentioned [CCS] result about the asymptotic normality of the h-index empirically, that immediately implies the concentration-about-the-mean property, (what Yong erroneously thought was only a conjecture, but was in fact a theorem).

Symbolic Moment Calculus

In [Z1] (see also [Z2] and [CJZ]) we initiated a symbolic-computational method for the automatic (and rigorous!) proof of limit laws for many families of combinatorial random variables. But with much lesser effort, one can always derive the same results *empirically* by a combination of *number-crunching* and *symbol-crunching* using the very *naive* approach that we will briefly recall.

Let X_n be an infinite sequence of combinatorial families (for example, $\{0,1\}^n$), and let $f(x)$ be a random variable (for example, the sum of the entries, alias the number of 1's).

Define a sequence of polynomials, $C_n(t)$, in a variable t, by

$$C_n(t) := \sum_{x \in X_n} t^{f(x)} \quad ,$$

(in the above example $C_n(t) = (1+t)^n$), called the *combinatorial generating functions*. Under the *uniform distribution*, this turns into *probability generating functions*, by dividing by $C_n(1)$ (alias $|X_n|$)

$$P_n(t) := \frac{C_n(t)}{C_n(1)} \quad .$$

(in the above example $C_n(t) = (1+t)^n/2^n$), To get the expectation, $E_n(f)$, let's call it a_n, one simply computes $\frac{d}{dt} P_n(t)\Big|_{t=1}$ (in the above example $a_n = n/2$). To get the *centralized* version, one divides $P_n(t)$ by t^{a_n}, getting

$$Q_n(t) := \frac{P_n(t)}{t^{a_n}} \quad .$$

The *variance*, $m_2(n)$, is given by

$$m_2(n) = (t\frac{d}{dt})^2 Q_n(t)\Big|_{t=1} \quad ,$$

and the higher moments, $k \geq 3$, by

$$m_k(n) = (t\frac{d}{dt})^k Q_n(t)\Big|_{t=1} \quad .$$

Finally, the *standardized moments*, $\alpha_k(n)$, are given by

$$\alpha_k(n) = \frac{m_k(n)}{m_2(n)^{k/2}} \quad .$$

The random variable f has a *limiting distribution* if for every $k > 2$,

$$\beta_k := \lim_{n\to\infty} \alpha_k(n) \quad ,$$

exists. This is usually a *continuous* probability distribution, and **very often** happens to be the good old *normal distribution* $\frac{1}{\sqrt{2\pi}} e^{-x^2/2}$, whose moments are 0 for k odd, and $\frac{k!}{(k/2)!2^{k/2}}$ for k even. In that case our family of combinatorial random variables is called *asymptotically normal*.

In *many* cases, this approach can be used to teach the computer to prove **rigorous** results, by getting either closed-form, or recursive descriptions, of the leading asymptotics, in n, of the moments $m_k(n)$, for **symbolic** n and k, and from which one can easily get the leading asymptotics for the standardized moments, $\alpha_k(n)$. This happens when the $C_n(t)$ are either given explicitly, or via a decent *grand generating function*

$$F(q,t) := \sum_{n=0}^{\infty} C_n(t) q^n \quad ,$$

where $F(q,t)$ is more-or-less explicit.

But what if $F(q,t)$ is not so nice? Then we can abandon (alleged) 'rigor', and do things *empirically*. Use $F(q,t)$ to crank out the first 10000 or whatever, polynomials $C_n(t)$, and do all the above steps up to, say, the 14-th moment, and estimate asymptotics of $\alpha_k(n)$ for $k \leq 14$ or whatever. If the leading terms seem to agree with those of the (standard) normal distribution: $1, 0, 3, 0, 15, 0, 105, 0, 945, \ldots$ we have a very convincing *empirical proof* of asymptotic normality. Also, as a bonus we can numerically estimate asymptotic expressions for the average a_n, and the variance $m_2(n)$.

The Hirsch (formerly Durfee) Polynomials

Alexander Yong reminds us (Eq. 1 of [A]), about the famous **Euler-Gauss** identity

$$\prod_{i=1}^{\infty} \frac{1}{1-q^i} = \sum_{k=0}^{\infty} \frac{q^{k^2}}{\prod_{j=1}^{k}(1-q^j)^2} \quad .$$

This lovely identity was given a pretty combinatorial proof by the Sylvester school (Durfee was Sylvester's graduate student), that could be found, for example, in George Andrews' partition **bible**([A], pp. 27-28).

That proof immediately implies that the *grand generating function*

$$\sum_{n=0}^{\infty} C_n(t) q^n \quad ,$$

equals

$$\sum_{k=0}^{\infty} \frac{q^{k^2} t^k}{\prod_{j=1}^{k} (1-q^j)^2} \quad .$$

(Note that in order to get the first N^2 members of the sequence $C_n(t)$ we only need to take the sum up to $k = N$ and then `taylor` it up to q^{N^2}.)

This generating function is not so easy to handle, but Maple can be easily used to crank-out the first 10000 terms. The first 6400 members of the *probability generating functions* (i.e. $C_n(t)/C_n(1) = C_n(t)/p_n$) can be found in *Maple input format*, suitable for computer-experimentation, in the file

`http://www.math.rutgers.edu/~zeilberg/tokhniot/oHIRSCH1` .

The first 100 terms of the subsequence consisting of perfect squares, i.e. the list of $C_{i^2}(t)$, for $1 \leq i \leq 100$ can be found here:

`http://www.math.rutgers.edu/~zeilberg/tokhniot/oHIRSCH2` .

The statistical analysis, as outlined above, can be found in the file

`http://www.math.rutgers.edu/~zeilberg/tokhniot/oHIRSCH3` .

Let's summarize our findings (**Warning**: these are non-rigorous, (but reliable), estimates).

$$a_n = 0.5404446395\,\sqrt{n} + 0.085691 + 0.0374788\,\frac{1}{\sqrt{n}} + O(\frac{1}{n}) \quad .$$

$$m_2(n) = 0.081057\,\sqrt{n} + 0.018459 - 0.018015\,\frac{1}{\sqrt{n}} + O(\frac{1}{n}) \quad .$$

Finally for the fourth through the eighth standardized even moments, we have:

$$\alpha_4(n) = 3.000000000 - .084847493\,\frac{1}{\sqrt{n}} - 0.1071813\,\frac{1}{n} + O(\frac{1}{n^{3/2}}) \quad ,$$

$$\alpha_6(n) = 15.0000000 - 12.60947794\,\frac{1}{\sqrt{n}} + 2.080133651\,\frac{1}{2} + O(\frac{1}{n^{3/2}}) \quad ,$$

$$\alpha_8(n) = 105.0000000 - 174.8856\,\frac{1}{\sqrt{n}} + 104.0909\,\frac{1}{n} + O(\frac{1}{n^{3/2}}) \quad ,$$

etc. etc. **Very convincing!**.

As already noted in the abstract, the variance is proportional (asymptotically) to the average. It follows that, since the h-index is asymptotically normal, that there is **concentration about the mean**.

The Maple Package HIRSCH

Readers are welcome to continue to explore, and generalize, by downloading the Maple package HIRSCH, already mentioned in the abstract, whose url is:

`http://www.math.rutgers.edu/~zeilberg/tokhniot/HIRSCH` .

References

[A] G. Andrews, *"The Theory of Partitions"*, Cambridge University Press, 1998. Originally published by Addison-Wesley, 1976.

[CCS] E.R. Canfield, S. Corteel and C.D. Savage, *Durfee polynomials*, Elec. J. Comb. **5** (1998), R32. (21 pages). Available on-line from:
`http://www.combinatorics.org/ojs/index.php/eljc/article/view/v5i1r32/pdf` .

[CJZ] E.R. Canfield, S. Janson, and D. Zeilberger, *The Mahonian Probability Distribution on Words is Asymptotically Normal*, Advances in Applied Mathematics **46** (2011), 109-124. [Special issue in honor of Dennis Stanton]. Available on-line from:
`http://www.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/mahon.html` .

[H] J. E. Hirsch, *An index to quantify an individual's scientific research output*, Proceedings of the National Academy of Science, **102 # 46** (Sept. 1, 2005), 16569-16572. Available on-line from:
`http://www.pnas.org/content/102/46/16569.full` .

[Y] A. Yong, *Critique of Hirsch's Citation Index: a combinatorial Fermi problem*, Notices of the American Mathematical Society **61 #9** (Oct. 2014), 1040-1050. Available on-line from:
`www.ams.org/notices/201409/rnoti-p1040.pdf` .

[Z1] D. Zeilberger, *The Automatic Central Limit Theorems Generator (and Much More!)*, "Advances in Combinatorial Mathematics: Proceedings of the Waterloo Workshop in Computer Algebra 2008 in honor of Georgy P. Egorychev", chapter 8, pp. 165-174, (I.Kotsireas, E.Zima, eds. Springer Verlag, 2009.) Available on-line from:
`http://www.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/georgy.html` .

[Z2] D. Zeilberger, *HISTABRUT: A Maple Package for Symbol-Crunching in Probability theory*, The Personal Journal of Shalosh B. Ekhad and Doron Zeilberger, Aug. 25, 2010. Available on-line from:
`http://www.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/histabrut.html` .

Doron Zeilberger, Department of Mathematics, Rutgers University (New Brunswick), Hill Center-Busch Campus, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA.
zeilberg at math dot rutgers edu ; http://www.math.rutgers.edu/~zeilberg/ .

Shalosh B. Ekhad, c/o D. Zeilberger, Department of Mathematics, Rutgers University (New Brunswick), Hill Center-Busch Campus, 110 Frelinghuysen Rd., Piscataway, NJ 08854-8019, USA.

EXCLUSIVELY PUBLISHED IN : http://www.math.rutgers.edu/~zeilberg/pj.html and arxiv.org.

Oct. 31, 2014